

In Acute Gastric Variceal Bleeding & Endoscopic Variceal Obliteration

Rx

ENDOCRYL®

n-Butyl Cyanoacrylate 0.5ml & 1ml


FAST & Reliable Tissue Adhesive



Making India Proud since decades



Edge Over Band Ligation







Variceal Bleeding



Polymerizes in 5 to 10 seconds after contact with blood²

Recommended in the case of bleeding from³



Endocryl Kit

- n-Butyl Cyanoacrylate 1.0 ml — 1 ampoule
- Instrument Cleansing Fluid 2.0 ml — 1 bottle
- Sclerotherapy Needle 21 G — 1 unit
- Syringe 2 ml — 2 units

Reference: 1. Tundak M et al., Ann Hepatol. 2013; 13(1): 75-83. 2. Desai P.N., VideoGIE, April 2021; 6(4): 155-158.
3. Kozial S et al., Prz Gastroenterol. 2015; 10(4): 239-243. IGV-Isolated Gastric Varix GOV-Gastroesophageal varices
NBICA- n-Butyl Cyanoacrylate

India's Pioneer
TISSUE ADHESIVE



END O CRYL[®]

Now in **0.25 ml**

For Topical Wound Closure

In Topical Wound Closure

 **ENDOCRYL[®]**

n-Butyl Cyanoacrylate 0.25ml

Simpler, Smoother & Safer

Wound Closure

Stronger & Tighter bond
within **15 secs¹**

Sloughs off naturally
within **5-10 days¹**

Forms **anti-microbial
barrier** over the wound²



ALICIA

A Division of
SAMARTH[®]
LIFE SCIENCES PVT. LTD.

Samarth House, 106, Bangor Nagar, off Link Road, Near Ayappa Temple, Coimbatore (W), Marathi - 400 008
• Tel.: 92235218017 / 02235218043 • Email: info@samarthlife.com • www.samarthlife.com

¹ www.spicell.com
² Pageer AJ, et al. Am J Surg. 2004; 148: 695-698

For the use of a Registered Medical Practitioner at a Hospital or Laboratory only.